

Updates on Contraceptive Technology Part 2

Kirsten Vogelsong
Mario Festin
Promoting Family Planning
Department of Reproductive Health and Research




Training Course in Sexual and Reproductive Health Research
Geneva 2016

08_XXX_MM1

Comparing Effectiveness of Family Planning Methods

More effective

Less than 1 pregnancy per 100 women in one year




How to make your method more effective

Implants, IUD, female sterilization:

After procedure, little or nothing to do or remember

Vasectomy: Use another method for first 3 months




Injectables: Get repeat injections on time

Lactational Amenorrhea Method (for 6 months): Breastfeed often, day and night


Pills: Take a pill at the same time each day

Patch, ring: Keep in place, change on time



Condoms, diaphragm: Use correctly every time you have sex

Fertility awareness methods: Abstain or use condoms on fertile days. Standard Days Method and Two-Day Method may be easier to use.



Withdrawal, spermicides: Use correctly every time you have sex

Less effective

About 30 pregnancies per 100 women in one year

Outline and Objectives

- Description of the method
- Mechanism of action
- Effectiveness
- Eligibility criteria
- Benefits and side effects
- Interventions for associated effects

08_XXX_MM3

Methods

- Combined oral contraceptives
- Progestin only pills
- Injectable contraceptives, progestin-only
- Injectable contraceptives, combined
- Hormonal implants
- **IUDs (copper bearing)**
- **LNG IUS**
- **Male and female condoms**
- **Other barrier methods**
- **Fertility awareness, lactational amenorrhea**
- **Emergency contraception**
- **Tubal ligation and vasectomy**
- **Other methods**


Copper Intrauterine Contraceptive Devices (IUDs or IUCDs)



Copper T 380A

08_XXX_MM5


What is a Copper IUD?



- Small plastic device with copper sleeves or wire around it, inserted through the vagina and cervix into the uterus
 - Safe
 - Highly effective
 - Long acting (up to 12 years)
 - Require trained provider to insert and remove

08_XXX_MM6

Copper-bearing IUD



Copper T-380A

TCu-380A, “Copper T” is most widely used copper IUD




Multiload 375

Multiload 375

Copper T: Mechanism of Action

Prevents fertilization by:

- Impairing the viability of the sperm
- Interfering with movement of the sperm



Source: Ortiz, 1996

Copper IUDs: Characteristics

- Safe and highly effective
- Require no user action
- Long-acting (up to 12 years)
- Rapid return to fertility
- No systemic effects
- Other health benefits
- Potential side effects
- Require pelvic exam
- Trained provider needed to insert and remove
- Possible pain or discomfort during insertion
- Do not protect against STIs/HIV
- Rare complications

Source: CCP and WHO, 2011

Copper IUDs: Possible Side Effects

Some women may experience:

- More cramps and pain during menstruation
- Increased or prolonged menstrual bleeding
- Bleeding between menstrual periods



Side effects are most common during the first 3–6 months.

Source: CCP and WHO, 2011; Larsson, 1993; DeMaeyer, 1989; WHO, 2004, updated 2008; WHO Special Programme of Research Development and Research Training in Human Reproduction, 1997

Copper IUDs: Health Benefits

IUDs are known to:

- Reduce risk of ectopic pregnancy
 - Rate in all IUD users is 12 in 10,000 (2 in 10,000 for Copper T380A users)
 - Rate in women using no contraception is 65 in 10,000
- Help protect against endometrial cancer

Source: CCP and WHO, 2011; Sivin, 1991.

Potential Complications

- Perforations—Very rare, 1 in 1,000 insertions
 - Linked to skill and experience of provider
 - Reduced through supervised training
- PID—Rare, most due to gonorrhoea or chlamydia at time of insertion
- Expulsions—Related to provider skill, age and parity of woman and insertion factors

08_XXX_MM12

Expulsion Rates Are Higher for Postpartum Insertion

Timing of Insertion	Expulsion Rates
Interval (more than 6 weeks after delivery)	Low (3% for skilled provider)
Immediate postpartum (within 10 minutes)	Slightly higher
Early postpartum (between 10 minutes and 48 hours)	Moderately higher
Late Postpartum (48 hours to 4 weeks)	High - Generally not recommended

Source: Chi, et al, 1985.

Who Can Initiate Copper IUDs

WHO MEC category 1 and 2 examples

WHO MEC Category	Conditions (selected examples)
Category 1	≥ 20 years, cervical ectopy, uterine fibroids without distortion of the uterine cavity, irregular bleeding without heavy bleeding, breastfeeding > 6 weeks, history of PID
Category 2	Menarche to < 20 years, nulliparous, heavy or prolonged bleeding, severe dysmenorrhea, anemia, HIV/AIDS infection, with or without ARV therapy

Source: WHO, 2004; updated 2008.

Who Should Not Initiate Copper IUDs

WHO MEC category 3 and 4 examples

WHO MEC Category	Conditions (selected examples)
Category 3	48 hours to <4 weeks postpartum, ovarian cancer/if initiating use, high individual risk of STI/ if initiating use
Category 4	pregnancy, unexplained vaginal bleeding (prior to evaluation), current PID or cervical infection, endometrial or cervical cancer/if initiating use

Source: WHO, 2004; updated 2008.

Timing of IUD Insertion

Interval insertion

- Within the first 12 days of menstrual cycle
- Any other time if woman is not pregnant

Postpartum insertion

- Immediately after vaginal or cesarean delivery if no infection or bleeding complications
- Within 48 hours or delay at least 4 weeks

Postabortion insertion

- Immediately or within 12 days if no infection

Source: WHO, 2004; updated 2008.

Side Effects

- Common side effects:
 - Heavier and/or prolonged menstrual bleeding
 - Menstrual cramping
 - Spotting between periods
- Side effects:
 - Are not signs of illness
 - Usually become less within the first 3–6 months

Correcting Misconceptions

IUDs:

- Rarely lead to PID
- Do not increase risk of STIs, including HIV
- Do not work by causing abortion
- Do not make women infertile
- Do not move to the heart or brain
- Do not cause birth defects
- Do not cause pain for either partner during sex
- Significantly reduce risk of ectopic pregnancy

08_XXX_MM18



Signs of Possible IUD Complications

- Bleeding and severe abdominal cramping within a few days post-insertion ➡ perforation
- Irregular bleeding or pain every cycle ➡ partial expulsion, perforation
- Fever, unusual vaginal discharge, low abdominal pain ➡ infection
- Missing IUD strings, missed period ➡ expulsion, pregnancy

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications:
Heavy, Prolonged or Irregular Bleeding

Counseling and reassurance are key

Problem	Action/Management
Heavy or prolonged bleeding	<ul style="list-style-type: none">● Reassure client that this is common and not harmful, usually diminishes after few months● For short-term relief offer 5-day course of tranexamic acid or NSAIDs (not aspirin)● Provide iron tablets
Irregular bleeding	<ul style="list-style-type: none">● Reassure client● For short-term relief offer ibuprofen or indomethacin 2 times daily after meals for 5 days

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications: Cramping and Mild Pain

Counseling and reassurance are key

Problem	Action/Management
Cramping and mild pain	<ul style="list-style-type: none">● She can expect cramping and pain in first 1–2 days after insertion● Reassure client that this is common in first 3–6 months, not harmful, usually decreases over time● Suggest ibuprofen, other pain reliever (not aspirin if she also has heavy bleeding)● If cramping continues, occurs outside of menstruation, evaluate, treat or refer● If cramping is severe but no underlying condition, discuss removing the IUD

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications: Severe Pain in Lower Abdomen (Rare)

- Rule out PID, ectopic pregnancy or perforation.
- If PID is suspected, treat with appropriate antibiotics for gonorrhoea, chlamydia and anaerobic bacterial infection. There is no need to remove the IUD.
- If ectopic pregnancy is suspected, refer immediately.

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications: Suspected Perforation

- Stop procedure immediately, remove IUD
- Observe vital signs for an hour; check for signs of bleeding
 - If rapid pulse, falling blood pressure, or increased pain: refer
- Provide alternative contraception
 - Advise avoid sex for 2 weeks
- Follow-up in a week or as needed

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications: Missing Strings

- Determine risk of pregnancy
- Perform pelvic exam, probe for strings in cervical canal
- If cannot locate strings, consider X-ray or ultrasound, or refer
- Give choice of another contraceptive method
- Insert another IUD if expulsion is confirmed and
 - Woman is not pregnant
 - She still wants to use an IUD

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications:

Suspected Pregnancy

- Assess for pregnancy, including ectopic pregnancy
- If the client is pregnant and wishes to continue the pregnancy:
 - Explain that using an IUD during pregnancy increases the risk of preterm delivery or miscarriage
 - If possible, remove the IUD
 - If not possible to remove, advise close follow-up for signs of septic miscarriage

Source: CCP and WHO, 2011.

Hormonal intrauterine device



LNG-IUS

08_XXX_MM26

Hormonal IUDs




- Steadily release a progestin
- Levonorgestrel IUD (LNG-IUD) most common hormonal IUD
- Also known as LNG-intrauterine system (LNG-IUS)
- Immediately reversible
- Effective for 5 years

Other features of the LNG IUS

- Works primarily by suppressing the growth of the lining of the uterus
- Needs to be inserted into a uterus by a trained health care provider
- Some report lighter, fewer or even absent bleeding days
- Also infrequent reports of headaches, breast tenderness or pain, acne

08_XXX_MM28


Male and Female Condoms



08_XXX_MM29

What Is a Male Condom?

- A sheath that fits over a man's erect penis
- Most are made of thin latex rubber
- Other materials include natural skin and various synthetics



08_XXX_MM30

What Is a Female Condom?

- A transparent sheath that fits loosely inside a woman's vagina
- Most common type is FC2—made of thin, soft, synthetic rubber film
 - Has flexible rings at both ends
- Natural latex female condoms available in some countries



Comparing Effectiveness of Male and Female Condoms

In real-life situations, correct and consistent use may be difficult to achieve

Pregnancy rates:

Male



Female




perfect use	2%	5%
typical use	15%	21%

Source: Hatcher, 2007.


Condoms: Mechanism of Action

- **Create a barrier that keeps sperm from entering the vagina, thus preventing pregnancy**
- **Also form a barrier against STIs including HIV**



08_XXX_MM33

Male and Female Condoms
Dual Protection



Condoms are the *only* FP method that provides **dual protection**:

- Protection from pregnancy *and*
- Protection from transmission of HIV and other STIs between partners

Effectiveness for Preventing HIV and STIs

- The consistent, correct use of condoms significantly reduces the risk of HIV infection in men and women
- When used correctly with every act of sex, condoms are 80%–95% effective in preventing HIV infection
- Condoms prevent STIs transmitted through body fluids such as gonorrhea and chlamydia
- Condoms are less effective for preventing STIs transmitted by skin-to-skin contact, such as herpes and warts

Characteristics of Male Condoms



- Safe and easy to use
- Widely available
- Effective when used consistently and correctly
- Provide dual protection
- No hormonal side effects
- Can help men with premature ejaculation
- Do not require provider's help
- Can be used as temporary backup method
- Protect women from conditions caused by STIs

08_XXX_MM36


Characteristics of Male Condoms

- As typically used, less effective than many other methods
- Require partner communication and cooperation
- Can be damaged by oil-based lubricants, heat, humidity or light
- May reduce sensation
- Can interrupt sex



Who Can Use Male Condoms

- All men and women can safely use male condoms, *except* those with severe allergic reaction to latex
 - Extremely rare among both men and women
 - Non-latex condoms are available in some countries



08_XXX_MM38

Male Condoms

Managing Problems

- If condom not used—Offer emergency contraception
- If slipping or breaking—Ask about practices, behaviors
- If difficulty putting on—Ask client to demonstrate, correct errors
- Difficulty persuading partner to use—Help her choose approaches that will work
Suggest adding another method



Male Condoms

Managing Problems

Mild irritation or mild allergic reaction to condom—Itching, redness, rash and/or swelling

- Try another brand of condoms
- Put lubricant or water on the condom
- If symptoms continue, assess or refer for possible vaginal infection or STI
- If no infection, may have allergy to latex

Characteristics of **Female Condoms**

What Men and Women Like

What Women Like:

- Female-controlled
- Texture feels more natural than latex male condoms
- Offer STI/HIV protection
- Outer ring provides stimulation
- Do not require provider's help

What Men Like:

- Can be inserted in advance
- Are not tight or constricting
- Do not dull sensation
- Do not have to be removed immediately

08_XXX_MM41

Characteristics of **Female Condoms**

What Men and Women Don't Like


Limitations:

- May be difficult to insert at first, require practice
- Not as effective as other methods
- More expensive than male condoms
- Less available than male condoms

08_XXX_MM42

Who Can Use Female Condoms


- All men and women can use FC2
- Latex female condoms: Severe allergic reaction to latex is the only condition that prevents use
 - Extremely rare



08_XXX_MM43

Managing Problems

- If having trouble inserting: Ask her to demonstrate
- If uncomfortable: Reinsert or reposition
- If condom squeaks: Use more lubricant
- If condoms slips or is not used correctly: Offer emergency contraception
- Difficulty persuading partner: Help her choose approaches that will work



Female Condoms

Managing Problems

(Continued)

- Mild irritation in or around the vagina or penis (itching, redness, rash)
 - Usually goes away on its own
 - Suggest added lubricant inside condom or on penis
 - If symptoms persist, assess and treat for possible vaginal infection or STI

08_XXX_MM45

Responding to Myths and Fears

Male condoms:

- Do not make men impotent, weak, or decrease sex drive
- Do not cause illness in men because sperm “backs-up”

Female condoms:

- Are not difficult to use, but correct use needs to be learned
- Do not cause illness in women because they prevent semen from entering the body

Responding to Myths and Fears

(continued)

Both male and female condoms:

- Cannot get lost in woman's body
- Do not have holes that HIV can pass through
- Are not intended only for use outside of marriage, but also used by married couples

08_XXX_MM47

Other barrier methods

- **Diaphragm** – is placed deep in the vagina before sex, covering the cervix as a barrier, usually used with a spermicide
- When used correctly with every act of sex, 6 pregnancies per 100 women using the diaphragm over the first year.
- **Cervical caps** – is also placed deep in the vagina before sex, covering the cervix as a barrier.
- Not as effective, with 20 pregnancies per 100 women in the first year.



Other barrier methods

- Spermicide – applied deep in the vagina before every act of sex
 - Usually Nonoxynol 9, but other substances also available
 - Work by causing the membrane of sperm cells to break, killing them or slowing movement.
 - One of least effective methods, may be a primary or a back up method.
- Microbicide (either anti-infective alone or dual protection)



Fertility awareness methods

- Requires the cooperation of both partners, with a commitment to abstain or use another method on fertile days
 - Periodic abstinence
 - Natural family planning
- Must be aware of the body changes or keep track of days, according to rules of the specific methods.
- No side effects.

08_XXX_MM50

Fertility awareness methods

- Calendar based methods
 - Standard days methods
 - Calendar rhythm methods
- Symptoms based methods
 - Cervical secretions
 - Basal Body Temperature
 - Increase noted slightly after release of the egg or ovulation
 - Two day method
 - Sympto-thermal method


08_XXX_MM51

Standard Days Methods

- 95% effective with consistent and correct use
- A woman can use the SDM if most of her menstrual periods are 26 to 32 days long.
- Days 8 to 19 of each cycle are considered as fertile days.
- The couple can use cycle beads, a color-coded string of beads that indicates fertile and nonfertile days.



How to use cycle beads



- Move ring to RED bead when period starts

1



- Move ring to next bead every day. Move ring even on bleeding days

2



- Use condoms or abstain when ring is on WHITE beads

3




- BROWN beads are safe days of no pregnancy

4



- When period starts again move ring to red bead to begin again.

5



- Always check your period comes between dark brown bead and last brown bead.

Lactational amenorrhea

- A family planning method based on breastfeeding
- Can be effective for up to 6 months after delivery, as long as monthly bleeding has not returned and the woman is fully breastfeeding.
- Provides an opportunity to offer a woman an ongoing method for continuously 6 months

08_XXX_MM54

Lactational amenorrhea

- A woman with the following may want to consider other methods:
 - Has HIV infection
 - Especially if the woman is not taking ARVs, not fully breastfeeding, newly infected or has advanced disease.
 - Using certain medications (mood-altering drugs, reserpine, ergotamine, anti-metabolites, cyclosporine, high dose corticosteroids, bromocriptine, radioactive drugs, lithium and certain anticoagulants)
 - A newborn with a condition that makes it difficult to breastfeed

08_XXX_MM55

Emergency contraception

- Emergency contraceptive pills help to prevent pregnancy, if taken up to 5 days after unprotected sex and other situations such as
 - forced sex (rape) and contraceptive mistakes (condom slippage or breakage, misplaced IUD, late for injections, etc)
- Do not disrupt an existing pregnancy
- Safe for all women
- Provide an opportunity for women to start using an ongoing FP method
- Effectiveness (no method = 8 pregnancies/100 woman years)
 - Progestin only ECPs – 1 pregnancy per 100 women years
 - Combined estrogen progestin ECPs – 2 pregnancies per 100 women years.
- Copper IUD effective as emergency contraception, when inserted up to 7 days after unprotected sex

Emergency contraception

- Contains a progestin alone (levonorgestrel or ulipristal) or a progestin and an estrogen together
- Works primarily by preventing or delaying ovulation
- Does not work if the woman is already pregnant.

08_XXX_MM57

Pill type	Total dosage to provide
Levonorgestrel-only dedicated product	<ul style="list-style-type: none"> • 1.5 mg of levonorgestrel in a single dose.⁶
Estrogen-progestin dedicated product	<ul style="list-style-type: none"> • 0.1 mg ethinyl estradiol + 0.5 mg levonorgestrel. Follow with same dose 12 hours later.
Progestin-only pills with levonorgestrel or norgestrel	<ul style="list-style-type: none"> • Levonorgestrel pills: 1.5 mg levonorgestrel in a single dose. • Norgestrel pills: 3 mg norgestrel in a single dose.
Combined (estrogen-progestin) oral contraceptives containing levonorgestrel, norgestrel, or norethindrone	<ul style="list-style-type: none"> • Estrogen and levonorgestrel pills: 0.1 mg ethinyl estradiol + 0.5 mg levonorgestrel. Follow with same dose 12 hours later. • Estrogen and norgestrel pills: 0.1 mg ethinyl estradiol + 1 mg norgestrel. Follow with same dose 12 hours later. • Estrogen and norethindrone pills: 0.1 mg ethinyl estradiol + 2 mg norethindrone. Follow with same dose 12 hours later.

Correcting misconceptions about Emergency contraception


ECPs :

- Do not cause abortion
- Do not cause birth defects if pregnancy occurs
- Are not dangerous to a woman's health
- Do not promote sexual risk taking
- Do not make women infertile

08_XXX_MM59


Female Sterilization Tubal Ligation

- Permanent method, done through mini-laparotomy (small incision) or laparoscopy
- One of the most effective methods with less than 5 failures per 1000 women
- May be used by any woman
- Counselling needed to ensure no post procedural regrets



Male Sterilization

Vasectomy



- Simple permanent surgical method in men
- With a 3 month delay in taking effect due to storage of sperm
- Tubes that carry sperm to the penis (vas deferens) are cut and ligated (by sutures or cautery)
- May be done using non-scalpel technique
- Less than 2 per 1000 pregnancies over first year
- Does not affect male performance nor provides increase risk for cancer

Other methods

- Ring

- Requires keeping a flexible ring which releases progestins and estrogen which are absorbed by the body
- Kept for 3 weeks, and woman menstruates in 4th week
- Prevents ovulation

- Patch

- Requires wearing a small adhesive patch with estrogen and progestins weekly, for 3 weeks
- Works by preventing ovulation



Acknowledgments and References

- Main Reference:
 - Family Planning – A Global Handbook for Providers (<https://www.fphandbook.org/>)
- Acknowledgements
 - Family Health International
 - Knowledge for Health
 - Institute of Reproductive Health