



ANTENATAL, INTRAPARTUM &
POSTNATAL CARE

DEFINITIONS

- **Antenatal care (ANC):** Care provided *during pregnancy* by skilled health-care professionals to ensure best health conditions for both mother and baby.
- **Intrapartum care:** Care provided *during childbirth* by skilled health-care professionals to ensure best health conditions for both mother and baby.
- **Postnatal care (PNC):** Care provided *up to six weeks following childbirth* by skilled health-care professionals to ensure best health conditions for both mother and baby.


RATIONALE – 1/2

- **In many contexts, adolescent pregnancy is common:** In 2016, an estimated 21 million girls aged 15-19 in developing countries became pregnant, approximately 12 million of whom gave birth. An estimated 2.5 million girls aged under 16 years in low-resource countries give birth every year. Drivers are context specific; they include child marriage, poverty, lack of opportunity & values related to womanhood & motherhood.
- **Adverse maternal health outcomes among adolescents have major health & social consequences:** Pregnancy & childbirth complications are the leading cause of death among girls aged 15-19 years globally. In addition, adolescent mothers face higher risks of maternal morbidity. Early child bearing can increase risks for newborns as well as for young mothers.

RATIONALE – 1/2

- **ANC, IPC & PNC are effective:** The health benefits of these interventions for mother and baby are clear. There are not ascertainable harms or burdens.
- **Access to & provision of good quality services needs attention:** Adolescents face barriers to accessing & using skilled care before, during & after pregnancy. Certain groups of adolescents e.g. very young adolescents, unmarried adolescents, & those who are displaced because of war, civil strife or other emergencies face special barriers.


Percentage of adolescent girls aged 15–19 years attended by a service provider at least four times during pregnancy (ANC 4), 2013–2018



Source: UNICEF's SOWC 2019 - data analyzed by International Center for Equity in Health, Federal University of Pelotas, Brazil, based on DHS, MICS and other national surveys, 2019.

Note: *Data refer to the most recent year available during the period specified in the chart title. Regional estimates represent data from countries representing at least 50 per cent of the regional population. Data coverage was insufficient to calculate regional estimates for East Asia and the Pacific, Europe and Central Asia, Middle East and North Africa, Latin America and the Caribbean and North America. The boundaries shown and the designations used on this map do not imply official endorsement or acceptance by the United Nations.

Percentage of births among adolescent mothers aged 15–19 years attended by skilled health personnel (typically a doctor, nurse or midwife), by country, 2013–2018



Source: Joint UNICEF/WHO SBA database, based on DHS, MICS and other national surveys as well as national administrative data, 2019.

Note: *Data refer to the most recent year available during the period specified in the chart title. Regional estimates represent data from countries representing at least 50 per cent of the regional population. Data coverage was insufficient to calculate regional estimates for East Asia and the Pacific, Europe and Central Asia, Latin America and the Caribbean, and North America. The boundaries shown and the designations used on this map do not imply official endorsement or acceptance by the United Nations.



HUMAN RIGHTS OBLIGATIONS

- States are obliged under human rights law to provide ANC, IPC & PNC.
- Upholding adolescents' rights in this area is linked to state obligations to ensure universal access to a comprehensive package of SRH interventions before, during & after pregnancy to all women & girls.
- Maternal health care should be free, confidential, adolescent-responsive and non-discriminatory; third-party authorization requirements should be removed.

KEY CONCEPTS TO CONSIDER

- **Pregnant adolescents, especially unmarried ones, often face barriers to accessing maternal health services including ANC, IPC & PNC:** Ensure availability of & access to ANC, IPC & PNC, including emergency obstetric care.
- **ANC, IPC & PNC services are often not responsive to the needs of adolescents:** It is critical for health workers to receive pre- & in-service training, & ongoing support to ensure they have the competencies & attitudes to provide high quality care, based on the rights of all people to health, confidentiality & non-discrimination.

WHO GUIDELINES

- *WHO guidelines on preventing early pregnancy and poor reproductive outcomes among adolescents in developing countries (2011).*
- *WHO recommendations on antenatal care for a positive pregnancy experience (2016) – with an update on micronutrient supplementation (date to be added).*
- *Use of multiple micronutrient powders for point-of-use fortification of foods consumed by pregnant women (2016).*
- *Optimal serum and red blood cell folate concentrations in women of reproductive age for prevention of neural tube effects (2015).*
- *Guidelines for the identification and management of substance use and substance use disorders in pregnancy (2014).*
- *WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia (2011) - with updates in 2018 and 2020)*
- *WHO recommendations for induction of labour (2011 – with updates on induction of labour at our beyond term, 2018).*
- *WHO recommendations for augmentation of labour (2014).*
- *WHO recommendations for intrapartum care for a positive childbirth experience (2018).*
- *WHO recommendations for prevention and treatment of maternal peripartum infections (2015).*
- *WHO recommendations for the prevention and treatment of postpartum haemorrhage (2012) – with updates in 2018 and 2020.*
- *WHO recommendations on tranexamic acid for the treatment of postpartum haemorrhage (2017).*
- *Daily iron supplementation in postpartum women: guideline (2016).*
- *Optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting (2012).*
- *WHO recommendations on health promotion interventions for maternal and newborn health (2015).*
- *Optimizing health worker roles to improve access to key maternal and newborn health interventions through task shifting (2012). WHO recommendations on health promotion interventions for maternal and newborn health (2015).*

COMPLEMENTARY GUIDELINES TO WHO'S GUIDELINES

- **Pregnancy, childbirth, postpartum and newborn care: a guide for essential practice, 3rd edn. (WHO, 2015).**
- **Managing complications in pregnancy and childbirth: a guide for midwives and doctors, 2nd edn, (WHO, 2017).**
- **Companion of choice during labour and childbirth for improved quality of care: evidence-to-action brief (WHO, 2016) – updated version dated 2020.**
- **Prevention and elimination of disrespect and abuse during childbirth (WHO, 2014).**
- **Obstetric fistula: guiding principles for clinical management and programme development (WHO, 2006).**
- **WHO recommendation on duration of bladder catheterization after surgical repair of simple obstetric urinary fistula (WHO, 2020)**
- **Not on pause: Responding to the sexual and reproductive health needs of adolescents in the context of the COVID-19 crisis (UNFPA, 2020)**
- **ASRH Tool Kit for Humanitarian Settings (Inter-agency working group, 2020)**

4

PROVISION OF ANTENATAL, INTRAPARTUM AND POSTNATAL CARE




RECOMMENDED ACTION – Maintain and modify

SPECIFIC MEASURES FOR DELIVERY OF SERVICES

- Inform adolescents where and how to access maternal care through mass media and digital media where adolescents have access to them.
- Consider using telemedicine for counselling and screening, including for risk factors known to be increased in the context of COVID-19 and to which adolescents may be particularly vulnerable (e.g. mental health conditions and gender-based violence) and the occurrence of danger signs.
- Where comprehensive facility-based services are disrupted, (i) prioritize antenatal care contacts for pregnant adolescents, (ii) ensure that birth preparedness and complication readiness plans are adapted at each contact to consider changes to services and (iii) prioritize postnatal care contacts during the first week after childbirth.
- Put in place targeted outreach strategies where coverage and care-seeking among pregnant adolescents have declined.

For further recommendations see [UNFPA Technical Brief package on facility-based maternity service delivery and phone-based antenatal and postnatal care during the COVID-19 pandemic](#).





**The pregnancy is progressing well.
Everything is in order. Now I
would like to discuss with you about
planning and preparing for the birth**

**Thank you for raising this. We
are worried about this because
of the night curfews**

ANTENATAL, INTRAPARTUM, POSTNATAL CARE

A REGIONAL PERSPECTIVE



a call for
solidarity
and action




World Health
Organization

REGIONAL OFFICE FOR THE Eastern Mediterranean

Maternal Mortality Ratio/ 100,000 live births

- **EMR** has the second highest MMR globally
- **Somalia** has the highest MMR (829/100,000 LB)
- **Afghanistan** is next (638/100,000 LB)¹




a call for
solidarity
and action







World Health
Organization

REGIONAL OFFICE FOR THE Eastern Mediterranean

Maternal conditions as leading cause of mortality among female adolescent 15-19 years ²



-  Countries with maternal conditions as number one cause of death
-  Countries with maternal conditions as number two cause of death
-  Countries with maternal conditions as number three cause of death
-  Countries where maternal conditions are not among the top 5 causes of death

Regional- Key indicators

- **Maternal healthcare indicators**

- I. **Antenatal care coverage at least four visits (%)**

- Somalia has the least coverage (6%). UAE has the highest proportion of women with antenatal coverage (97%).³

- II. **Proportion of births attended by skilled birth attendance (SBA)**

- SBA in Afghanistan (59%) and Somalia (9.4%)⁴

- III. **Proportion of mothers who had postnatal contact with a health provider within 2 days of delivery**

- Somalia and Yemen are the only two countries in the Region that have less than 20% of postnatal care who had postnatal contact with a health provider.⁵

- **Legal and policy support for maternal health services**

- I. 15/16 surveyed countries have **national policies on antenatal and postnatal care for mother and newborn.**⁶

- II. 15/16 countries have the **policy regarding the right of every women to access skilled birth attendance.**⁷

- III. Somalia is the only country that does not have policy for the provision these health services.

Barriers at regional level to accessing health care for pregnant adolescents ⁸

■ Individual factors

- I. Limited autonomy (decision making power).
- II. Low literacy rate among adolescent girls.
- III. Lack of awareness and information about their health and existing health services.
- IV. Financial constraints.

■ Socio-cultural factors


- I. Deep rooted social norms around gender and sexuality/gender discrimination.
- II. Child marriage / teenage pregnancy.
- III. Perception of community and service providers towards unmarried and/or unregistered married girls (judgmental attitude).
- IV. Socio-economical determinants, overprotection of girls.

■ Health services provision


- I. Data measurement and shortage of age disaggregated data/ Birth registration.
- II. Lack of availability of services: MCH services often do not focus on young first-time mothers.
- III. Lack of access and quality of care: Poor infrastructures, privacy, and confidentiality, competent health worker/ cost of services.
- IV. Disrespect and abuse during childbirth.⁹
- V. Conflicts, humanitarian settings, Covid-19 pandemic.

Effect of COVID-19 on coverage of maternal health services in selected countries in EMR ¹⁰


Sudan: % of women delivered by skilled birth attendant




Sudan: Number of women who attended PNC2



Yemen: Number of ANC visits/contacts provided



Pakistan: Number of facility births



Addressing regional challenges with Safe Motherhood initiative ^{8,11}

- **Delay marriage and first birth:**

- I. Investment in girl's education, employment and economical dependency.
- II. Judicial and law enforcement infrastructure on minimum age of marriage law which should be publicized and enforced in the local context with supporting, associated policies in place.
- III. Ensure access of adolescents and youth to information on sexual and reproductive health education within and out of school context.
- IV. Increase availability and access for family planning services with focus on reducing unmet need among young people.
- V. Community mobilization and engagement to support interventions to prevent child marriage.

- **Increasing access to SRH including maternal health services:**

- I. Establish and strengthen integrated safe motherhood programme at various levels of care with focus on unreachable young mothers. Task shifting / Community based interventions / Mobile and outreach services.
- II. Advocating for improved SRH/MNH policy and practice for adolescent girls & young mothers.
- III. Comprehensive programmatic approach to address mental health support and violence prevention.
- IV. Improving the quality of services by training the healthcare workers to address the needs of adolescent mother in a friendly environment.

Regional initiative 1: Advocacy campaign in Pakistan ¹²

In Feb 2012 CARE (Leading humanitarian and development organization) Pakistan started the project titled:

“Advocating for improved maternal newborn health (MNH) and sexual reproductive health (SRH) policy and practice for adolescent girls and young mothers (AIMS)”

The 14 months project was Implemented in partnership with Rahnuma-Family Planning Association of Pakistan (FPAP).

- **Aim**
 - To increase awareness regarding the specific reproductive and sexual health needs of adolescent girls and young mothers.
 - To advocate for their inclusion in provincial health policies in four provinces of Pakistan.
- **Outcomes**
 - A robust advocacy strategy was designed.
 - Advocacy campaign was planned for Federal and provincial levels.
 - Various key stakeholders of MNH & SRH were willing to accept their responsibilities for ensuring a supportive environment for youth.

Regional initiative 2: Improving maternal and neonatal health in Sudan Oct 2011 - Dec 2012 ^{13,14}

A community-based intervention project implemented in Kassala Town and Rural Kassala localities, covering the catchment communities of 19 PHCUs.

- **Objectives**


Strengthening Primary Health Care and Community Mobilization to contribute to improved health and wellbeing of mothers, newborns and their families.

- **Outcomes**

- Enhanced the capacity of the PHUs with training of staff and providing supplies.
- The introduction of the referral fund to cover the cost of the antenatal and other maternity care services.
- Community mobilization through establishment of networks and channels to reach discrete audiences at their residential areas through women as an “agent” of change.
- Continuity of supply of family planning methods.

Regional opportunities

Regional alignment with Global Initiatives/Guidance



- **Global strategy for women, children, and adolescents** ¹⁵

It is an opportunity for the countries to integrate ASRH interventions to achieve the global target of reducing MMR to less than 70/100,000 LB (SDG 3.1). The plan is to strengthen the healthcare delivery system, improve skills of health workers on early detection and management of complications in pregnancy and delivery.¹¹

- **Adolescent Health in All Policies (AHiAP)** ¹⁶

It is a strategy to consider the implications of decisions on adolescent health, avoid harmful effects and seek synergies. It facilitates the formulation of adolescent-responsive public policies in sectors other than health.

- **Vision 2023 Health for All by All** ¹⁷

Universal health coverage: inclusion of ASRH evidence-based interventions in the regional priority benefit package.¹³

References

1. WHO Regional Office for the Eastern Mediterranean. Monitoring health and health system performance in the Eastern Mediterranean Region: Core indicators and indicators on the health-related Sustainable Development Goals 2019. Cairo: WHO Regional Office for the Eastern Mediterranean; 2020. Licence: CC BY-NC-SA 3.0 IGO. Available from: https://rho.emro.who.int/sites/default/files/booklets/EMR-HIS-and-core-indicators-2019-final_0.pdf%20
2. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: Adolescent country profile. World Health Organization; c2021. Available from: <https://www.who.int/data/maternal-newborn-child-adolescent-ageing/static-visualizations/adolescent-country-profile>
3. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: Antenatal care coverage - at least four visits (%). World Health Organization; c2021. Available from: [https://www.who.int/data/maternal-newborn-child-adolescent-ageing/indicator-explorer-new/mca/antenatal-care-coverage---at-least-four-visits-\(-\)](https://www.who.int/data/maternal-newborn-child-adolescent-ageing/indicator-explorer-new/mca/antenatal-care-coverage---at-least-four-visits-(-))
4. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: Proportion of births attended by skilled health personnel (SDG 3.1.2). World Health Organization; c2021. Available from: <https://www.who.int/data/maternal-newborn-child-adolescent-ageing/indicator-explorer-new/mca/proportion-of-births-attended-by-skilled-health-personnel>
5. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: Proportion of mothers who had postnatal contact with a health provider within 2 days of delivery. World Health Organization; c2021. Available from: <https://www.who.int/data/maternal-newborn-child-adolescent-ageing/indicator-explorer-new/mca/proportion-of-mothers-who-had-postnatal-contact-with-a-health-provider-within-2-days-of-delivery>
6. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: National policy on postnatal care for mothers and newborns. World Health Organization; c2021. Available from: <https://www.who.int/data/maternal-newborn-child-adolescent-ageing/national-policies/mca/national-policy-on-postnatal-care-for-mothers-and-newborns?themelD=3af0211e-8037-4fad-b36d-60614c4b37b4>
7. WHO Global Observatory database. Maternal, newborn, child and adolescent health and ageing data portal: National policy/guideline on right of every women to have access to skilled care at birth. World Health Organization; c2021. Available from: <https://www.who.int/data/maternal-newborn-child-adolescent-ageing/national-policies/mca/national-policy-guideline-on-right-of-every-women-to-have-access-to-skilled-care-at-childbirth?themelD=3af0211e-8037-4fad-b36d-60614c4b37b4>

References

8. Mahaini R. Improving maternal health to achieve the Millennium Development Goals in the Eastern Mediterranean Region: a youth lens. East Mediterr Health J. 2008;14 Suppl:S97-106.
9. WHO. WHO statement on prevention and elimination of disrespect and abuse during facility-based childbirth. World Health Organization; 2015. Available from: https://apps.who.int/iris/bitstream/handle/10665/134588/WHO_RHR_14.23_eng.pdf;jsessionid=B7B5554C4B229F0A00CDD01C3882BE98?sequence=1
10. WHO EMRO Country dashboard of monitoring the effect COVID-19 on SRMNCAH. (unpublished data).
11. Safe Motherhood. Priorities for safe motherhood. Safe Motherhood; c2016. Available from: <http://www.safemotherhood.org/priorities/index.html>
12. CARE International in Pakistan. Advocating for improved MNH/SRH Policy and Practice for adolescent girls and young mothers: Key achievements, lessons learnt and way forward recommendations. CARE International in Pakistan; 2013. Available from: https://www.care.org/wp-content/uploads/2020/05/PSJ-2013-Way_Forward_Report.pdf
13. Improving maternal and neonatal health in Kassala State: strengthening primary health care and community mobilization in Kassala town and rural Kassala localities. Oct 2011-Dec 2012. (unpublished).
14. Abdelrahman Mohamed Mubarak. Evaluation report of a project on: Improving maternal and neonatal health in Kassala State - Strengthening primary health care & community mobilization in Kassala town and rural Kassala localities. (unpublished).
15. Every Woman Every Child. Global Strategy for Women's, Children's and Adolescents Health 2016-2030. Every Woman Every Child; 2015. Available from: <https://www.who.int/life-course/publications/global-strategy-2016-2030/en/>
16. WHO. Global Accelerated Action for the Health of Adolescents (AA-HA!): guidance to support country implementation. Geneva: World Health Organization; 2017. Licence: CC BY-NC-SA 3.0 IGO. Available from: <https://apps.who.int/iris/bitstream/handle/10665/255415/9789241512343-eng.pdf?sequence=1>
17. WHO Regional Office for the Eastern Mediterranean. Vision 2023 for Eastern Mediterranean Region. World Health Organization; c2021. Available from: <http://www.emro.who.int/about-who/vision2023/vision-2023.html>